

# Identification of complement factor 5 as a susceptibility locus for experimental allergic asthma

Christopher L. Karp<sup>1,\*</sup>, Andrew Grupe<sup>2,\*</sup>, Eric Schadt<sup>2</sup>, Susan L. Ewart<sup>3</sup>, Michelle Keane-Moore<sup>4</sup>, Peter J. Cuomo<sup>4</sup>, Jörg Köhl<sup>5</sup>, Larry Wahl<sup>6</sup>, Douglas Kuperman<sup>4</sup>, Soren Germer<sup>7</sup>, Dee Aud<sup>2</sup>, Gary Peltz<sup>2</sup> and Marsha Wills-Karp<sup>4</sup>

The prevalence and severity of allergic asthma continue to rise, lending urgency to the search for environmental triggers and genetic substrates. Using microarray analysis of pulmonary gene expression and single nucleotide polymorphism-based genotyping, combined with quantitative trait locus analysis, we identified the gene encoding complement factor 5 (C5) as a susceptibility locus for allergen-induced airway hyperresponsiveness in a murine model of asthma. A deletion in the coding sequence of C5 leads to C5-deficiency and susceptibility. Interleukin 12 (IL-12) is able to prevent or reverse experimental allergic asthma. Blockade of the C5a receptor rendered human monocytes unable to produce IL-12, mimicking blunted IL-12 production by macrophages from C5-deficient mice and providing a mechanism for the regulation of susceptibility to asthma by C5. The role of complement in modulating susceptibility to asthma highlights the importance of immunoregulatory events at the interface of innate and adaptive immunity in disease pathogenesis.

The worldwide prevalence and severity of allergic asthma have increased dramatically in recent decades. Unfortunately therapeutic advances have not kept pace, and asthma morbidity and mortality continue to rise<sup>1</sup>. The cardinal features of allergic asthma include airway hyperresponsiveness (AHR) to a variety of specific and nonspecific stimuli, excessive airway mucus production, pulmonary eosinophilia and elevated concentrations of serum immunoglobulin E (IgE). It is generally accepted that asthma arises as a result of inappropriate immunological responses to common environmental antigens in genetically susceptible individuals<sup>2</sup>. Pathophysiology is thought to be mediated by CD4<sup>+</sup> T lymphocytes producing a type 2 cytokine profile<sup>3</sup>. The molecular mechanisms that underlie susceptibility to these aberrant immune responses are unknown. A genetic approach has promised insight into mechanism. Family studies demonstrate a heritable predisposition to asthma. The mapping of asthma susceptibility genes in humans has been hampered by variability in phenotype, genetic heterogeneity across populations and uncontrolled environmental influences. Despite this, more than 20 linkage regions have been found<sup>4,5</sup>, although the path from linkage to gene remains unraveled to date.

A well characterized murine model that mimics the pathophysiology of human allergic asthma was used in a reductive approach to these complexities<sup>6</sup>. In this model, allergen exposure results in AHR,

increased airway mucus content, antigen-specific IgE in serum, and pulmonary eosinophilia<sup>3,7</sup>. As with human asthma, the pulmonary inflammatory response in experimental asthma centers around the airway wall and involves extensive airway wall remodeling, including: goblet cell metaplasia; thickening of the airway epithelial layer; and airway smooth muscle hypertrophy. Inbred mouse strains vary in their susceptibility to disease induction in this model. Allergic airway





Figure 1. C5 gene expression and genotype correlate with allergen-induced AHR. C5 transcript levels in whole lungs of OVA-sensitized A/J (□), C5H/HeJ (○), A/J × C5H/HeJ (●), 6 BC1-high responder and 6 BC1-low responder mice are shown in comparison with their dynamic airway pressure (APTI in cm-H<sub>2</sub>O × s). BC1-mice carrying only the A/J C5 allele (■); BC1-mice that are C5 heterozygous (□). The outlier BC1 mouse (homozygous for the A/J C5 allele, low APTI) is discussed in the text. Expression units reflect the average difference value as analyzed by GeneChip 3.01 Software.

<sup>1</sup>Department of Medicine, Johns Hopkins University, School of Medicine, Baltimore, MD, USA. <sup>2</sup>Roche BioScience, 3401 Hillview Avenue, Palo Alto, CA 94304-1397, USA.

<sup>3</sup>Department of Large Animal Sciences, Michigan State University, B41 National Food Safety and Toxicology, East Lansing, MI 48824-1302, USA. <sup>4</sup>Department of Environmental Health Sciences, Johns Hopkins University School of Hygiene and Public Health, Baltimore, 615 N. Wolfe St., Baltimore, MD 21205, USA. <sup>5</sup>Institute für Medical Microbiology, Medizinische Hochschule Hannover, Carl-Neuberg-Str. 1, 30625 Hannover, Germany. <sup>6</sup>National Institute of Dental and Craniofacial Research, 30 Convent Drive, Bethesda, MD 20892 USA. <sup>7</sup>Roche Molecular Systems, Inc., 1145 Avenue of the Americas, Alameda, CA 94501, USA. \*These authors contributed equally to this work.

Correspondence should be addressed to M.W.K. (mkarp@welch.jhu.edu).



**Figure 2.** C5 allele correlates with allergen-induced AHR across inbred strains. A/J have a 5' deletion in the C5 gene and carry the C5-D allele; C5H/HeJ mice do not have the deletion and carry the C5-S allele. Allele-specific kinetic, polyacrylamide gel electrophoresis (PCR) was used to genotype other inbred mouse strains. (a) Kinetic PCR using the C5H/HeJ allele-specific primer identifies the C5-S allele in three strains. Primer dimer formation is detected in strain A/J. (b) Dynamic airway pressure (APTI) in congenitally sensitized mice.

lacking this allele after amplification cycle 38. (b) Kinetic PCR using the A/J allele-specific primer identifies the C5-D in two strains. (c) Dynamic airway pressure (APTI) in congenitally sensitized mice.

responses are mediated by cytokines produced by type 2 CD4<sup>+</sup> T cells in susceptible strains of mice. Congruent with the well defined role of Interleukin 12 (IL-12) in promoting type 1 CD4<sup>+</sup> T cell differentiation, disease is prevented or ablated by exogenous IL-12 in susceptible strains<sup>24</sup>. Similarly, neutralization of IL-12 renders resistant strains susceptible to disease<sup>25</sup>. Previous studies have shown that the various features of the asthmatic phenotype are separable, both genetically and experimentally in this model. Two strains with markedly different susceptibilities to experimental allergen-induced AHR, the primary phenotypic signature of human asthma, were used in a search for genetic substrates.

## Results

### C5 as a candidate gene


To identify gene candidates, pulmonary gene expression was profiled with oligonucleotide microarrays. A/J (highly susceptible to allergen-induced AHR) and C5H/HeJ (highly resistant) mice were immunized intraperitoneally and were subsequently challenged intratracheally with soluble ovalbumin (OVA); the allergic phenotype was assessed 3 days after the antigen challenge. After phenotypic assessment, lungs were collected from A/J, C5H/HeJ (A/J × C5H/HeJ)<sub>F1</sub> (called hereafter F<sub>1</sub>) and eight of the segregating F<sub>1</sub> × A/J backcross (BC<sub>1</sub>) mice that exhibited phenotypically extreme allergen-induced airway responsiveness (APTI). (APTI of BC<sub>1</sub>-high, 1280–2349 cm-H<sub>2</sub>O × s; APTI of BC<sub>1</sub>-low, 256–514 cm-H<sub>2</sub>O × s.) Of the 7350 genes on the microarray, 2718 were expressed in the lungs of the parental strains and 739 were differentially expressed in the lungs of A/J and C5H/HeJ mice. Approximately 227 genes exhibited a greater than threefold change in expression when these two parental strains were compared.

Differential gene expression was assessed within groups of high- and low-responder mice by comparing expression in four BC<sub>1</sub>-high responder and A/J mice, and four BC<sub>1</sub>-low responder and C5H/HeJ mice, respectively. The expression data was filtered for differential expression in four of the five intragroup pairwise comparisons, as suggested

by the Affymetrix GeneChip software. This analysis yielded 21 differentially expressed genes. Previous analysis of the inheritance pattern of allergen-induced AHR in these strains led to the identification of two distinct quantitative trait loci (QTL) on chromosome 2: *Abhr1* ("allergen-induced bronchial hyperresponsiveness") and *Abhr2* (lod scores=4.3 and 3.7, respectively). C5, located near *Abhr2* at 23.5 cM on chromosome 2, was the only gene that both met the differential expression criteria and was located in one of the defined QTL intervals. No gene within the *Abhr1* support interval satisfied the differential expression criteria.


The initial expression data set was confirmed by expression profiling of additional BC<sub>1</sub>-high responder (*n*=2), BC<sub>1</sub>-low responder (*n*=2) and parental (A/J and C5H/HeJ) mice, to yield a total of 18 data points. Direct comparison of allergen-induced AHR with pulmonary C5 mRNA expression in A/J, C5H/HeJ, F<sub>1</sub> and 12 BC<sub>1</sub> mice exhibiting extreme allergen-induced AHR phenotypes revealed that resistant (C5H/HeJ and BC<sub>1</sub>-low responder) mice had high C5 gene expression; susceptible (A/J and BC<sub>1</sub>-high responder) mice had low C5 gene expression; and F<sub>1</sub> mice had intermediate C5 gene expression (Fig. 1). The amount of C5 expression was significantly associated with genotype (*P*<0.005 for the BC<sub>1</sub> mice alone; *P*<0.001 including the parental and F<sub>1</sub> mice as well, with Student's *t*-test).

C5 expression also correlated with the magnitude of allergen-induced AHR, linear correlation coefficient (*r*<sup>2</sup>) = -0.66. The C5 genotype was



**Figure 3.** Effects of C5aR blockade on cytokine production by primary human monocytes. Monocytes were incubated in serum-free conditions with the indicated concentration of C5aR antagonist (\*) or control (○). Stimulation was with IFN-γ + SAC (solid line) or SAC alone (dashed line). (a) IL-12 p70 production. (b) IL-12 p40 production. (c) TNF-α production. (d) IL-10 production. Values shown are means ± s.d. in a representative experiment (*n*=4). Statistical analysis was done with paired *t*-tests. \**P*<0.005, \*\**P*<0.01, \*\*\**P*<0.005. At 1 μM of the C5aR antagonist, there was 95.4% (s.e. 3%) mean inhibition of IL-12 p70 production; 81.7% (s.e. 6.3%) inhibition of IL-12 p40 production to stimulation with IFN-γ + SAC, 91.6% (s.e. 4.6%) to SAC alone; 63.3% (s.e. 5.7%) inhibition of TNF-α production to IFN-γ + SAC, 51.3% (s.e. 6.1%) to SAC alone; and 25.4% (s.e. 6.8%) mean augmentation in IL-10 production after stimulation with IFN-γ + SAC, -0.4% (s.e. 14.7%) after SAC alone (*n*=4).

**Figure 4. C5aR signaling and C5aR blockade have mechanistically distinct effects on IL-12 production.** Serum dependence and pertussis toxin sensitivity of signaling. Monocytes were incubated in media without (filled bars) or with (open bars) 250 ng/ml of pertussis toxin in the presence or absence of 10% fetal bovine serum for 2 h. Cells were subsequently cultured (with or without serum) in the presence of 100 nM C5aR antagonist or 100 nM recombinant C5a (Sigma), or appropriate controls (control lysate<sup>†</sup> for C5aR, media for C5a). Stimulation was with IFN- $\gamma$  + SAC. Data presented are means  $\pm$  s.d. in a representative experiment ( $n=3$ ). Statistical analysis was done by paired *t*-test. \* $P<0.000007$ ,  $\dagger P=0.000004$ ,  $\ddagger P=0.00008$ .



analyzed for all BC<sub>1</sub> mice shown in Fig. 1. Five out of six BC<sub>1</sub>-low responder mice were homozygous at the C3 locus, and had high pulmonary C5 mRNA expression. The single BC<sub>1</sub>-low responder mouse with low pulmonary C5 mRNA expression was homozygous for the A/J allele at the C3 locus, a result that is congruent with the QTL analysis demonstrating the presence of at least two loci controlling allergen-induced AHR in this model.

### SNP-based genotypic analysis

The relationship between C3 expression and genotype has been examined previously in murine systems. A/J mice have a 2-bp deletion in a 5' exon of the C3 gene that renders them deficient in C3 mRNA and protein production (and devoid of functional C3); C3H/He mice are C3-sufficient<sup>11</sup>. The link between C3 and AHR was explored further by characterizing the correlation between C3 genotype and susceptibility to allergen-induced AHR. A high-throughput single nucleotide polymorphism (SNP)-based genotyping assay, employing C3 allele-specific oligonucleotide primers and kinetic thermal cycling, was used to genotype 172 BC<sub>1</sub> mice with phenotypically extreme allergen-induced AHR responses.

BC<sub>1</sub>-high responder mice tended to be homozygous for A/J C3 alleles (C3-D); whereas BC<sub>1</sub>-low responder mice tended to be heterozygous, carrying both C3-D and C3H/He (C3-S) alleles (Table 1). These differences in C3 allele frequency were statistically significant, confirming the correlation between C3 genotype and susceptibility to allergen-induced AHR in the progeny of these parental strains. The correlation between phenotype and genotype at this locus is not absolute, confirming the presence of multigenic influences on allergen-induced AHR. As a control, these same BC<sub>1</sub> mice underwent SNP-based genotyping of the Toll-like receptor 4 gene (*Tlr4*) on chromosome 4, a dominant negative mutation in which is responsible for endotoxin hyporesponsiveness in C3H/He mice<sup>12,13</sup>. No significant difference in *Tlr4* allele frequency was found between the high- and low-responder populations (Table 1).

To begin to assess the breadth of the linkage between C3 genotype and susceptibility to allergen-induced AHR, SNP-based genotypic analysis was done in other, previously phenotyped, inbred murine strains (Fig. 2). The correlation held fast in this limited sample of strains. Both of the susceptible strains (AKR/J and A/J) carry the deletion in C3, whereas resistant (C57BL/6J, BALB/cJ, C3H/HeJ) strains do not carry this deletion.

### C5a effects on IL-12 production

C5 cleavage fragments exert pleiotropic effects on inflammatory responses. C5a has been shown to stimulate the production of proinflammatory cytokines by monocytes and macrophages. The effects of C5a on monocyte production of IL-12, a T helper cell subset 1 (Th1)-promoting proinflammatory cytokine that is critical to regulation of the asthmatic phenotype in this model, were therefore examined. *In vitro* analysis of the role of complement in modulation of monocyte and macrophage function is complicated, experimentally, by the facts that such cells synthesize the complement system autologously; complement activation occurs readily *in vitro*; and the use of serum provides access to complement activation fragments including C5a<sup>14-17</sup>. Therefore, to mimic the *in vivo* lack of access of A/J macrophages to C5a, we studied the effects of ablating C5a-mediated signaling in human monocytes cultured in the absence of exogenous complement. A potent, specific C5a receptor (C5aR) antagonist, isolated by panning C5a COOH-terminal libraries on a C5aR-expressing cell line, was used<sup>18</sup>. This antagonist binds to human leukocytes with an median inhibitory dose (ID<sub>50</sub>) in the low nanomolar range and exhibits no detectable agonist activity even at micromolar concentrations<sup>18</sup>.

Human monocytes were incubated under serum-free conditions with immunofluorescence-purified C5aR antagonist (or a control lysate similarly purified from bacteria not expressing recombinant C5aR antagonist), and secondarily stimulated for monokine production. Blockade of the C5aR caused marked, dose-dependent inhibition of the IL-12 productive capacity of primary human monocytes (Fig. 3a,b). This occurred both at the level of the functional p70 heterodimer and at the level of the highly regulated p40 subunit. K-76, a complement-inhibitory monocarboxylic acid derived from fungi with primary effects on C5 activation<sup>19</sup>, also inhibited IL-12 production by monocytes (data not shown). We examined the specificity of these effects by characterizing the effects of C5aR blockade on the principal pro- and anti-inflammatory cytokines produced by monocytes and macrophages: tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ) and IL-10. C5aR blockade also inhibited the secretion of TNF- $\alpha$ , albeit with less potency than its effects on IL-12 (Fig. 3c). Although inhibition of C5a-mediated signaling had no overall effect on bacterially driven IL-10 production, IFN- $\gamma$ -mediated suppression of IL-10 production was reversed (Fig. 3d). Taken together, these latter findings suggest that functional C3-deficiency leads to a relatively anti-inflammatory phenotype, especially in the presence of the type 1 cytokine IFN- $\gamma$ .


**Table 1. C3 genotype correlates with allergen-induced AHR in BC<sub>1</sub>-high responder and BC<sub>1</sub>-low responder mice**

alleles	Number of mice with allele type	
	BC <sub>1</sub> -high	BC <sub>1</sub> -low
<b>C3 genotype</b>		
AA <sup>a</sup>	54	32
AC <sup>a</sup>	32	54
<b>Th4 genotype</b>		
AA	36	46
AC	50	40

<sup>a</sup>AA, homozygous A/J allele; AC, heterozygous (A/J, C3H/He) alleles. Template DNA was prepared from 172 BC<sub>1</sub> mice exhibiting extreme phenotypes for allergen-induced AHR (APTI values for BC<sub>1</sub>-high, 1280–2348 cm-H<sub>2</sub>O  $\times$  s/APTI for BC<sub>1</sub>-low, 268–514 cm-H<sub>2</sub>O  $\times$  s). Genotyping was done by kinetic PCR with specific primers for A/J and C3H/He alleles of C3 and Th4. Statistical analysis was done using the Fisher exact test. C3 genotype:  $P=0.0013$ . Th4 genotype:  $P=0.169$ .

## ARTICLES

**Figure 6.** IL-12 production by macrophages from A/J and C3H/HeJ mice. Peritoneal macrophages from A/J (filled bars) and C3H/HeJ (open bars), isolated after thioglycolate elicitation, were stimulated with IFN- $\gamma$  + SAC, or mock-stimulated with media. After 24 h, IL-12 p70 production was measured by enzyme-linked immunosorbent assay (ELISA) (Pharmingen) in cell-free supernatants. Data presented are means  $\pm$  s.e. of 11 mice per strain. Statistical analysis was done by paired *t*-test. \**P* < 0.001.



It has recently been reported that C5a can itself lead to downmodulation of IL-12 production under certain conditions<sup>33,34</sup>. These data are replicable (Fig. 4). However the inhibition of IL-12 production induced by C5a is clearly and mechanistically separable from that resulting from ablation of C5aR signaling. The C5aR is coupled to a pertussis toxin-sensitive G protein<sup>35</sup>. Notably, C5a-mediated inhibition of IL-12 production is pertussis toxin sensitive and only occurs in the presence of serum. In contrast, C5aR antagonist-mediated inhibition is not pertussis toxin-sensitive and does not depend upon exogenous serum (Fig. 4).

To further investigate the relevance of these observations to experimental allergic asthma, IL-12 production was compared in A/J and C3H/HeJ mice. Peritoneal macrophages from C5-deficient A/J mice produce significantly less IL-12 than those obtained from C5-sufficient C3H/HeJ mice (Fig. 5).

## Discussion

We previously identified two QTL for allergen-induced AHR on chromosome 2, *Ahr1* and *Ahr2*<sup>28</sup>. To make the transition from QTL to gene, we took advantage of recent technological advances that have made genetic analysis of complex traits increasingly tractable in experimental models. QTL analysis was combined with microarray analysis of target organ gene expression and SNP-based genotyping to identify C5 as a susceptibility locus for AHR. Genotypic analysis of the phenotypically diverse backcross (BC<sub>1</sub>) mice was critical to investigation of the link discovered between phenotype, gene expression and genotype. The power of gene array analysis is tempered by its inability to distinguish between primary and secondary effects on gene expression. SNP analysis of the backcross animals demonstrated the presence of a genetic polymorphism at the C5 locus with a primary effect on the phenotype of allergen-induced AHR. The association identified between the C5 locus and AHR in our study was not complete, accurately reflecting the multigenic nature of this trait. As with any genetic analysis, it is theoretically possible that a closely linked gene or regulatory sequence<sup>36</sup>, and not the identified gene itself, underlies the trait in question. The role of C5 itself in allergen-induced AHR is supported by *in vitro* functional data on the effects of C5 cleavage fragment signaling on immunoregulatory cytokine production, however.

It has become clear in recent years that the complement system plays an important immunoregulatory role at the interface of innate and acquired immunity. For example, proteolytic fragments of C3 have been shown to influence the class of immune response to a given immunogen, augmenting humoral immune responses through effects on B cells, and suppressing cellular immune responses through inhibition of IL-12 production by antigen presenting cells<sup>37,38</sup>. Because experimental allergic asthma is clearly a cytokine-regulated process,

this suggested a mechanistic hypothesis for the role of C5-deficiency in susceptibility to allergen-induced AHR. Proteolytic cleavage of C5 yields two fragments, both of which can stimulate cytokine production. As part of a hemolytically inactive membrane attack complex, C5b causes signaling in neutrophils and endothelial cells, inducing chemokine production by the latter<sup>39,40</sup>. C5a has pleiotropic effects on inflammation, being chemotactic for all myeloid lineages, inducing degranulation and the production of variety of proinflammatory mediators by granulocytes and increasing vascular permeability<sup>41</sup>. C5a also stimulates monocyte and macrophage production of the proinflammatory cytokines TNF- $\alpha$ , IL-1 and IL-6<sup>42,43</sup>. We therefore examined the effects of C5a on monocyte production of IL-12. The data presented here demonstrate that blockade of C5aR signaling ablates IL-12 production by primary human monocytes. Such blockade also suppresses TNF- $\alpha$  production and reverses IFN- $\gamma$ -mediated suppression of IL-10 secretion, although it does not affect bacterially driven IL-10 production.

C5a has itself been shown to lead to IL-12 inhibition under certain conditions<sup>33,34</sup>, a finding that we were able to replicate. Our data demonstrate that C5a-mediated IL-12 suppression is mechanistically distinct from the IL-12 suppression that results from the ablation of C5aR signaling, however. These data, together with previously reported data showing a lack of agonist effects of the C5aR antagonist<sup>44</sup>, strongly suggest that the C5aR antagonist is not acting as an agonist at the receptor. Modulation of IL-12 by both the C5aR antagonist and C5a suggests a model in which some C5aR signaling is needed to render monocytes and macrophages competent for IL-12 production, whereas further exposure to C5a, especially in the presence of IFN- $\gamma$  generated during an ongoing inflammatory process, leads to inhibition of the production of this potentially toxic cytokine.

Defective IL-12 production by C5a-depleted monocytes and macrophages provides a plausible mechanism for the regulation of susceptibility to asthma by C5: IL-12 drives type 1 CD4<sup>+</sup> T cell responses, preventing or reversing experimental allergic asthma. Furthermore, ablation of C5-mediated signaling releases monocyte and macrophages from IFN- $\gamma$ -mediated inhibition of IL-10 production. Whereas IL-10 from antigen presenting cells can downmodulate both type 1- and type 2-polarized responses, feedback inhibition of IL-10 production by the type 1 cytokine IFN- $\gamma$  itself promotes ongoing type 1 responses. As a consequence of these effects on cytokine production by antigen presenting cells, the absence of C5 is permissive for type 2 responses, including allergic asthma. Its presence promotes type 1 responses.

As predicted by studies with human monocyte cells, macrophages from C5-deficient A/J mice produce significantly less IL-12 than macrophages from C5-sufficient C3H/HeJ mice. The effect of C5 genotype on maximal stimulation of IL-12 production by macrophages from these strains is not all-or-nothing. However, the differences in IL-12 productive capacity seen are biologically significant in the context of the *in vivo* development of an immunological response: IL-12 administration to A/J mice renders them resistant to the induction of asthma; neutralization of IL-12 in C3H/HeJ mice renders them susceptible<sup>4</sup>. Similar, biologically relevant, differences in IL-12 production by C5-deficient (A/J) and C5-sufficient (C57BL/6) strains have been found in a model of malaria infection<sup>45</sup>. Although a direct association between C5 deficiency and AHR in our model remains to be fully established, preliminary data from A/J mice in which the wild-type C5 gene has been restored suggest that the T<sub>H</sub>2-associated eosinophilic inflammatory response is attenuated in the presence of a functional C5 gene.

These results provide insight into other models of genetic deletion or antibody-mediated inhibition of C5, C5a, or the C5aR. Such models have suggested a central role for C5 in the pathogenesis of collagen-

induced arthritis, DTH responses, and endotoxic shock, as well as in resistance to *Listeria* and to blood-stage malaria infection<sup>23-26</sup>. Although C5a has pleiotropic effects, including important effects on leukocyte trafficking, all of these models are dependent on or exacerbated by IL-12, and downmodulated by IL-10<sup>27</sup>. Likewise, *in vivo* deficiencies of C5 and the C5aR are associated with blunted production of TNF- $\alpha$ <sup>28</sup>. Given the important role of complement at the interface of innate and adaptive immunity, it is likely that complement-associated genes will provide high quality candidates for susceptibility to other immune-mediated diseases.

Our identification of C5 as a susceptibility gene for experimental allergen-induced AHR may have relevance to human asthma. Two genome-wide screens for asthma susceptibility loci have found linkage to the C5 chromosomal region (9q34)<sup>29</sup>, and the human C5aR gene is located at 19q13.3, a chromosomal region with susceptibility loci for asthma in several different cohorts<sup>30</sup>. More generally, the likely role of C5-deficiency in suppressing the production of IL-12 in experimental asthma is echoed, pathophysiologically, by the finding that patients with allergic asthma have diminished production of IL-12 both in the lung and systemically<sup>31,32</sup>. Many of the candidate asthma susceptibility regions defined in human genetic studies contain genes whose products alter the balance of type 1 and type 2 cytokine expression. People with homozygous defects in the C5 gene, like others with deficiencies in late complement components, suffer a high incidence of disseminated necrotic infection. They are protected, however, from the fulminant shock that often accompanies such disease in the face of complement-susceptibility, a phenomenon that may be due to the effects of a lack of C5a on TNF- $\alpha$  and IL-12 production<sup>33</sup>. Whether C5 null relatives are prone to develop allergic asthma does not appear to have been addressed.

The work reported here shows the utility of combined genetic and genomic approaches to the analysis of complex traits in experimental rodent models. Although genetic susceptibility loci for human diseases may not be direct homologs of loci identified in this fashion, genes encoding other proteins in the implicated biological pathways are likely candidates. In any case, rapid identification of susceptibility loci in experimental animal models is likely to provide substantial insight into pathogenetic mechanisms in human diseases. Even in the absence of full genomic information, the power of the approach to susceptibility gene discovery outlined here is evident. With full sequencing of the human and murine genomes currently in sight, these methods should find broad usage in the search for genetic susceptibility loci underlying complex human diseases.

## Methods

**Induction and characterization of the allergic phenotype.** Four-week-old male mice (AJ, C5H/HeJ, AKR/J, BALB/c, C57BL/6J, and 4–6 mice per experimental group) and backcross mice (m172) were obtained from The Jackson Laboratory (Bar Harbor, ME) and housed under laminar flow hoods in an environmentally controlled specific pathogen-free animal facility for the duration of experiments. All mice were immunized by an intraperitoneal injection of 10  $\mu$ g OVA (Sigma) in 0.2 ml PBS buffer. Mice were anesthetized 14 days after immunization and challenged intratracheally with 50  $\mu$ l of a 1.5% solution of OVA. Characterization of the allergic phenotype was performed 72 h after antigen challenge. Mice were anesthetized, intubated, ventilated at a rate of 120 breaths per min with a constant tidal volume of air (0.2 ml), and paralyzed. After confirmation of a stable respiratory rate, 10  $\mu$ g per kg weight of acetylcholine was injected intravenously and dynamic airway pressure (APTI in cm-H<sub>2</sub>O  $\times$  s) was followed for 5 min.

**Gene array assays.** Gene chip arrays were hybridized with labeled cRNA obtained from whole lungs. Whole lungs, as opposed to cell type-specific antisense thereof, were employed to avoid bias: the relative importance to asthma pathogenesis of the multiple cell types that reside or traffic through the lung is unknown. Hybridizations were done with cRNA prepared from individual BALB/c mice; whereas lungs from AJ, C5H/HeJ and F<sub>1</sub> mice were obtained from four mice; lungs from m172 and C57BL/6J mice were obtained from four mice. Isolation of acetylcholine (2  $\times$  poly(A)<sup>+</sup>)-cRNA synthesis and hybridization of gene chips were performed as described in the *Experimental Procedures* section of the *Supplemental Material*.

**Technical Manual.** The image was obtained with a 4 $\times$  image axis. Differential expression, as analyzed by GeneChip 3.0 software from Affymetrix, Santa Clara, CA, was assessed by random pairwise comparisons of BC<sub>1</sub>-high and BC<sub>1</sub>-low responder mice, and by pairwise comparison of AJ and C5H/HeJ mice.

**SNP assays.** Genomic DNA was prepared from 176 phenotypically extreme BC mice. The APTI values for BC<sub>1</sub>-high responder mice were within the 90th percentile of APTI values from all BC mice. The APTI values of BC<sub>1</sub>-low responder mice were in the tenth percentile of all measured BC mice. DNA from inbred mouse strains was purchased from The Jackson Laboratory. Genotyping assays were performed by kinetic thermal cycling<sup>34</sup>. Allele-specific forward primers: C5-C5H/HeJ allele: AAGAATATCTTTTAAATTCAGAGTAT; C5-AJ allele: ACATATCTTTTAAATTCAGAGTAT; and a common reverse oligonucleotide primer: ACTATAGAGAGGATTTTACACCAAGCTGGAA. Tm4-C5H/HeJ allele: CCTTCACTACAGAGGATTTTATCTTC; Tm4-AJ allele: CCTTCACTACAGAGGATTTTATCTTC; and a common reverse oligonucleotide primer: CCTGGATGATTTTGGCAAGCA; were used to amplify the strain specific Tm4 alleles.

**In vitro assays of human monocyte function.** Human monocytes isolated by countercentrifugation from normal volunteers<sup>35</sup> were cultured adherently in Dulbecco's modified Eagle's medium (Gibco, Gaithersburg, MD) at a density of 2 $\times$ 10<sup>6</sup> cells (0.2 ml per data point) in 96-well plates (Costar, Cambridge, MA), in the presence or absence of varying concentrations of immunosilyl-purified, bacterially expressed recombinant C5aR antagonist, or a control preparation similarly purified from bacteria not expressing C5aR antagonist. Isolation and culture of monocytes were performed under lipopolysaccharide (LPS)-free conditions. After 2–3 h in culture, monocytes were stimulated with IFN- $\gamma$  (100 IU/ml; Pharmacia) followed 3 h later by lipopolysaccharide (LPS) from *Escherichia coli* (1:500; Calbiochem, La Jolla, CA), or with SAC alone. The production of p70 (as measured by p40) by cultured human monocytes is dependent upon preincubation with IFN- $\gamma$ . 24 h after SAC stimulation, cell-free culture supernatants were collected for measurement of cytokines by ELISA (IL-12 p40, IL-10, and TNF- $\alpha$  assays were from Pharmacia; IL-12 p70, from R&D Systems, Minneapolis).

All cell culture reagents were LPS-free to the limits of detection of the Limulus amoebocyte lysate assay (3–6 pg/ml) (BioWhittaker, Walkersville, MD). Bacterially modified lipopolysaccharide (BTL) was employed in studies using complement reagents with measurable LPS contamination. At 5  $\mu$ g/ml, the recombinant NH<sub>2</sub>-terminal modified fragment of BTL (BTL12, XOMA (US) LLC) has an LPS-neutralizing capacity of 210 ng/ml<sup>36</sup>. At experimental dilutions, all such reagents contain <100 ng/ml of LPS.

## Acknowledgements

Supported in part by NIH grants AI40507, DE12167 (C.L.K.), ES06066, HL58927 (M.W.-K.) and RR00097 (S.L.E.), BMRF grant 01VM9305 (L.K.), a Michigan State University Allergy Research Initiation Grant (S.L.E.), and EPA grant R625724 (M.W.-K.). The authors thank M. Shin for the gift of K-76.

Received 5 July 2000; accepted 3 August 2000.

- Swan, R. D. et al. National trends in the morbidity and mortality of asthma in the US. Prevalence, hospitalization and death from asthma over two decades: 1965–1994. *Chest* 91, 1053–1061 (1987).
- Wills-Karp, M. Immunologic basis of antigen-induced airway hyperresponsiveness. *Annu. Rev. Immunol.* 17, 255–281 (1999).
- Davies, S. E. et al. A genome-wide search for quantitative trait loci underlying asthma. *Nature* 383, 253–259 (1999).
- A genome-wide search for asthma susceptibility loci in isolically diverse populations: The Collaborative Study on the Genetics of Asthma (CSGA). *Nature Genet.* 16, 309–322 (1997).
- Deer, C. et al. A genome-wide search for asthma susceptibility loci in a founder population: The Collaborative Study on the Genetics of Asthma. *Hum. Mol. Genet.* 7, 1393–1398 (1998).
- Wijst, M. et al. A genome-wide search for linkage to asthma. *German Asthma Genetics Group. Genomics* 50, 1–3 (1999).
- Wills-Karp, M. et al. Interleukin-13: central mediator of allergic asthma. *Science* 282, 2258–2261 (1998).
- Gawell, H. et al. Interleukin-12 inhibits antigen-induced airway hyperresponsiveness, inflammation, and Th2 cytokine expression in mice. *J. Exp. Med.* 182, 1527–1536 (1995).
- Kane-McCoy, A., Wasyk, R., Finkbeiner, S. & Wills-Karp, M. Resistance to virus-induced airway hyperresponsiveness requires endogenous production of IL-12. *J. Immunol.* 161, 518–526 (1998).
- Ewart, S. E. et al. Identification of quantitative trait loci controlling airway hyperresponsiveness in inbred mice. *Am. J. Resp. Crit. Care Med.* 157, 1005–1012 (1998).
- Wills-Karp, M., Patterson, D. L. & Hershberg, D. L. Deficiency of the murine Th1-complement component (C3), A-2 gene pair gene deletion in a 5 $\times$  exon J. *Biol. Chem.* 268, 2431–2440 (1993).
- Poler, A. et al. Defective LPS signaling in C5H/HeJ and C57BL/6J mice: mutations in T4 gene. *Science* 282, 2095–2097 (1998).
- Vogel, S. H. et al. Cutting edge: functional characterization of the effect of the C5H/HeJ allele in mice that lack an *emg* gene: in vivo evidence for a dominant negative mutation. *J. Immunol.* 162, 5066–5069 (1999).
- Johnson, E. & Heston, G. Monoclonal: polyclonal: have the potential to synthesize the complete functional complement system. *Stand. J. Immunol.* 37, 489–493 (1990).
- Hidvegy, E. I. & Marks, R. M. Complement activation occurs on subendothelial extracellular matrix in vivo and is initiated by retention or removal of overlying endothelial cells. *J. Immunol.* 160, 3135–3138 (1998).
- Wasserman, K. O., Sayegh, G. & Blumberg, P. M. Macrophage type 3 complement receptors mediate serum-independent stimulation of Lactobacillus-derived dendritic cells and promote maturation of the parasite surface by immunotactile microscopy. *J. Exp. Med.* 184, 1332–1337 (1996).
- Berg, K. & Iversen, C. D. Production of monoclonal antibodies against the human angiotensin

- Cla des Arg and their application in the macrophage-specific sandwich-ELISA for the quantification of Cla des Arg in plasma. *J Immunol Methods* 152, 73-87 (1992).
18. Huhns T. et al. Selection of a C5a receptor antagonist from phage libraries attenuating the inflammatory response in immune complex disease and ischemic/reperfusion injury. *J Immunol* 163, 955-959 (1999).
  19. Hong S., Kikuchi T., Miyazaki M., Iwata T. & Iwata K. An anti-complement agent, K-76 monocarboxylic acid, its role and mechanism of inhibition of the complement activation cascade. *J Immunol* 122, 2418-2423 (1979).
  20. Wilmanns M. et al. C5a suppresses the production of IL-12 by IFN- $\gamma$ -primed and lipopolysaccharide-challenged human monocytes. *J Immunol* 162, 6763-6769 (1999).
  21. Braun M. C., Leiby E. & Kotewicz J. L. Selective suppression of IL-12 production by chemokines. *J Immunol* 164, 3009-3017 (2000).
  22. Garavito C. & Garavito N. P. C5a anaphylatoxin and its seven transmembrane-segment receptor. *Annu Rev Immunol* 12, 775-809 (1994).
  23. Lioy G. G. Identification of a coordinate regulator of interleukins 4, 13, and 5 by cross-species sequence comparisons. *Science* 288, 138-140 (2000).
  24. Fearon D. T. The complement system and adaptive immunity. *Smith Immunol* 10, 356-361 (1998).
  25. Mosser D. M. & Karg C. L. Receptor mediated subversion of macrophage cytokine production by intracellular pathogens. *Curr Opin Immunol* 11, 406-411 (1999).
  26. Wang C. et al. Homologous leucine C5a67 complex: an epitope of polymorphonuclear leukocytes. *Blood* 85, 2570-2578 (1995).
  27. Wang C., Garavito N. J. & Nicholson-Weller A. Signaling by homophilically inactive C5a67, an epitope of polymorphonuclear leukocytes. *J Immunol* 165, 785-792 (1995).
  28. Kikuchi T. S. et al. Sublytic concentrations of the membrane attack complex of complement induce endothelial interleukin-8 and monocyte chemotactic protein-1 through nuclear factor- $\kappa$ B activation. *Am J Pathol* 150, 2015-2021 (1997).
  29. Morgan E. L. et al. Identification and characterization of the effector region within human C5a responsible for stimulation of IL-6 synthesis. *J Immunol* 148, 3937-3942 (1992).
  30. Schneider R., Gellera J. A. & Chizzola C. A. Recombinant C5a stimulates transcription rather than translation of interleukin-1 (IL-1) and tumor necrosis factor: translation signal provided by lipopolysaccharide or IL-1 itself. *Blood* 78, 1631-1638 (1992).
  31. Cavatoni LM., Pilling C., Haefliger-Cavatoni N. Recombinant C5a enhances interleukin 1 and tumor necrosis factor release by lipopolysaccharide-stimulated monocytes and macrophages. *Eur J Immunol* 20, 263-267 (1990).
  32. Sano H. & Sawauchi M. H. Early IL-12 p70, but not p40, production by splenic macrophages correlates with host resistance to blood-stage *Plasmodium chabaudi* AS malaria. *Can. Exp. Immunol* 112, 343-349 (1998).
  33. Wang Y., Rotstein S. A., Madaï J. A. & Madaï L. A. Anti-C5 monoclonal antibody therapy prevents collagen-induced arthritis and ameliorates established disease. *Proc Natl Acad Sci USA* 92, 8955-8959 (1995).
  34. Topil R. E. et al. Required early complement activation in contact sensitivity with generation of local C5-dependent chemotactic activity and late T cell interferon  $\gamma$ : a possible initiating role of B cells. *J Exp Med* 180, 1015-1025 (1997).
  35. Garavito C., Samson M. & Samson E. Genetic control of resistance to *Leishmania* monocytes: regulation of leukocyte inflammatory responses by the Hc locus. *J Immunol* 132, 2078-2083 (1984).
  36. Barton P. A. & Warren J. S. Complement component C5 modulates the systemic tumor necrosis factor response in murine endotoxin shock. *Infect Immun* 61, 1474-1481 (1993).
  37. Tridandad G. Interleukin-12: a cytokine at the interface of inflammation and immunity. *Adv Immunol* 70, 83-243 (1998).
  38. Hogston U. L., Lu R., Gerard N. P. & Gerard C. Impaired inflammatory responses in the reverse primer reaction through genetic deletion of the C5a receptor. *J Exp Med* 188, 748-756 (1997).
  39. van der Pouw Kraan I. C. et al. Reduced production of IL-12 and IL-12 dependent IFN- $\gamma$  release in patients with allergic asthma. *J Immunol* 158, 5560-5565 (1997).
  40. Hasegawa T. et al. Expression of IL-12 and IL-13 mRNA in asthma and their regulation in response to steroid therapy. *Am J Respir Crit Care Med* 155, 845-851 (1997).
  41. Garavito C., Holland M. & Hignault R. High-throughput SNP allele frequency determination in pooled DNA samples by kinetic PCR. *Genome Res* 10, 260-268 (2000).
  42. Karg C. L. et al. Mechanism of suppression of cell-mediated immunity by measles virus. *Science* 272, 228-231 (1995).